ELSEVIER

Toxicology 116 (1997) 247–248

TOXICOLOGY

# Volume contents

## Volume 116 (1997)

The effect of age on aluminum retention in rats
**There are slight changes in the distribution of the minute fraction of aluminium absorbed from the gastrointestinal tract**
M. Gómez, D.J. Sánchez, J.M. Llobet, J. Corbella, J.L. Domingo (Spain) . . . . . 1–8

Cytokines (IL-1$\beta$ and TNF$\alpha$) in relation to biochemical and immunological effects of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) in rats
**TCDD elevates the mRNA levels of both cytokines**
F. Fan, B. Yan, G. Wood, M. Viluksela, K.K. Rozman (USA, Germany) . . . . . 9–16

Role of tachykinins and neutral endopeptidase in toluene diisocyanate-induced bronchial hyperresponsiveness in guinea pigs
**Tachykinin release accounts for this phenomenon as shown by capsaicin action**
F. Gagnaire, M. Ban, C. Cour, J.C. Micillino, P. Bonnet, D. Hettich (France) . . . . . 17–26

Enhancement of the stress response by minute amounts of cadmium in sensitized Reuber H35 hepatoma cells
**There is a biphasic change in sensitivity expressed as synthesis of heat shock proteins**
F.A.C. Wiegant, J. van Rijn, R. van Wijk (The Netherlands). . . . . 27–37

Long-term exposure of male and female mice to trivalent and hexavalent chromium compounds: effect on fertility
**12 weeks oral application decreases fertility in both sexes regardless of chromium valency**
A. Elbetieha, M.H. Al-Hamood (Jordan) . . . . . 39–47

Inhibition of DNA synthesis by paracetamol in different tissues of the rat in vivo
**At high doses there is a reversible inhibition of thymidine incorporation**
C.F. Lister, A.E.M. McLean (UK) . . . . . 49–57

Effects of arsenite pretreatment on the acute toxicity of parathion
**In Wistar rats, arsenite significantly lowers the $LD_{50}$ for parathion**
F.R. Siller, B. Quintanilla-Vega, M.E. Cebrián, A. Albores (México) . . . . . 59–65

Utilization of renal slices to evaluate the efficacy of chelating agents for removing mercury from the kidney
**Dimercaptosuccinic acid and meso-2, 3-dimercaptosuccinic acid are highly efficient chelators for mercury in kidney slices**
R.L. Keith, I. Setiarahardjo, Q. Fernando, H.V. Aposhian, A.J. Gandolfi (USA) . . . . . 67–75

Comparison of the toxicity profiles of ISIS 1082 and ISIS 2105, phosphorothioate oligonucleotides, following subacute intradermal administration in Sprague–Dawley rats
**Pathologic changes were seen in liver, kidney, spleen and bone marrow**
S.P. Henry, L.R. Grillone, J.L. Orr, R.H. Bruner, D.J. Kornbrust (USA) . . . . . 77–88

Sequence of exposure to cadmium and arsenic determines the extent of toxic effects in male Fischer rats
**A nontoxic dose of As followed by a toxic dose of Cd decreases Cd lethality, whereas Cd has no effect on As lethality in rats**
J.F. Hochadel, M.P. Waalkes (USA) . . . . . 89–98

Effects of dithiocarbamates on toxicity of cadmium in rat primary hepatocyte cultures
**Chelators appear to protect by lowering Cd levels in the cells, sequestrating reactive Cd ions and diminishing activity of reactive oxygen species**
T. Funakoshi, K.-i. Ueda, H. Shimada, S. Kojima (Japan) . . . . . 99–107

The acute effect of lead acetate on glucocorticoid receptor binding in C6 glioma cells
**Lead may perturb initial glucocorticoid binding events, possibly by affdcting PKC-mediated phosphorylation**
L.E. Tonner, D.I. Katz, A.S. Heiman (USA) . . . . . 109–122
Comparative effects of essential and nonessential metals on preimplantation mouse embryo development in vitro
**Essential metals examined were Cu, Mn, Fe and Zn and non-essential ones were Cr, Hg, Pb, V, Al, Ag, Cd and As**
L.A. Hanna, J.M. Peters, L.M. Wiley, M.S. Clegg, C.L. Keen (USA) . . . . . 123–131
Chronic exposure of rats to ozone and sulfuric acid aerosol: biochemical and structural responses
**There is no evidence for a synergistic interaction between ozone and sulfuric acid**
J.A. Last, K.E. Pinkerton (USA) . . . . . 133–146
Importance of cholinolytic drug selection for the efficacy of HI-6 against soman in rats
**Use of centrally acting cholinolytics causes very slight improvements over the peripheral actions of atropine**
J. Kassa (Czech Republic) . . . . . 147–152
Structure-activity relationships in the hydrolysis of acrylate and methacrylate esters by carboxylesterase in vitro
**The relative toxicity of acrylates does not appear to be due to differences in carboxylesterase-mediated hydrolysis**
T.J. McCarthy, G. Witz (USA) . . . . . 153–158
Prediction of teratogenic potency of valproate analogues using cerebellar aggregation cultures
**The proposed in vitro test yields values comparing well with the teratogenicity of valproate and its analogues**
T.E. Maar, U. Ellerbeck, E. Bock, H. Nau, A. Schousboe, V. Berezin (Denmark, Federal Republic of Germany) . 159–168
Selenite suppression of cadmium-induced testicular apoptosis
**Administration of selenite up to 2 hours before or after $CdCl_2$ is effective**
M.M. Jones, C. Xu, P. A. Ladd (USA) . . . . . 169–175
Protective effects of antioxidants against benomyl-induced lipid peroxidation and glutathione depletion in rats
**In vivo toxicity of benomyl may be associated with oxidative stress**
D. Banks, M.R.I. Soliman (USA) . . . . . 177–181
Testosterone pretreatment mitigates cadmium toxicity in male C57 mice but not in C3H mice
**Reduced lethality and hepatotoxicity may be due to testosterone enhancement of Cd-induced metallothionein synthesis**
H. Shimada, R.M. Bare, J.F. Hochadel, M.P. Waalkes (USA) . . . . . 183–191
Significant suppression of rat liver aldolase B by a toxic coplanar polychlorinated biphenyl, 3,3′,4,4′,5-pentachloro-biphenyl
**Activity was suppressed to about 50% of that measured in free- and pair-fed controls**
Y. Ishii, H. Kato, M. Hatsumura, T. Ishida, N. Ariyoshi, K. Oguri (Japan) . . . . . 193–199
Colchicine-induced elevation of tissue metallothionein contents is mediated by inflammation-independent serum factor
**Disruption of microtubules may release a mediator**
N. Itoh, M. Kasamatsu, S. Onosaka, N. Muto, K. Tanaka (Japan) . . . . . 201–209

**Immunotoxicology**
The effects of cyclosporin A, dexamethasone and other immunomodulatory drugs on induced expression of IL-3, IL-4 and IL-8 mRNA in a human mast cell line
**IL-3 mRNA expression in human leukemic mast cells is inhibited by cyclosporin and dexamethasone**
E.V. Warbrick, A.L. Thomas, C.M.M. Williams (UK) . . . . . 211–218
Immunomodulatory and protective effects of *N*-acetylcysteine in mitogen-activated murine splenocytes in vitro
***N*-acetylcysteine modulates the mitogen-activated blastogenesis of murine lymphocytes in vitro**
F.O. Omara, B.R. Blakley, J. Bernier, M. Fournier (Canada) . . . . . 219–226
Inhalation of diesel exhaust enhances antigen-specific IgE antibody production in mice
**Spleen cells of mice exposed to diesel exhaust inhalation were found to produce more IL-4 and IL-10, and less interferon-$\gamma$ than spleen cells from controls**
H. Fujimaki, K. Saneyoshi, F. Shiraishi, T. Imai, T. Endo (Japan) . . . . . 227–233

Announcements . . . . . 235–237

Author index . . . . . 239
Subject index . . . . . 241–245
Volume contents . . . . . 247–248